

DMOPC '15 Contest 1 P6 - Lelei and Contest

Time limit: 1.0s **Memory limit:** 256M

Lelei La Lalena has been studying competitive programming in our world. Today, she decides to do a contest on [DMOJ](#) to prove her skill! Confident, Lelei opens the sixth problem of the October 2015 DMOPC and finds a really abstract problem with no story. So she decides to make one up and tell **FatalEagle** to add it to the problem. Anyway, here's the original problem:

Rory is playing with an array A consisting of N integer elements indexed from 1 to N and a positive integer M . Rory will perform Q operations. Each operation is either type 1 or type 2.

Type 1 operation is in the form $1\ l\ r\ x$. You should add x to each element in $A[l], A[l+1], \dots, A[r]$.

Type 2 operation is in the form $2\ l\ r$. You should output the sum $(A[l]^M + A[l+1]^M + \dots + A[r]^M) \bmod M$.

Lelei is confident she can solve this problem, so she tells you that she doesn't need your help, as she can solve it faster than you. Seeing this as a challenge, you obviously want to show Lelei that she could have a better time penalty, if only she asked for your help. Can you prove her wrong?

Input Specification

The first line of input will contain three integers M , N , and Q .

The second line of input will contain N elements, the original elements of array A in the order $A[1], A[2], \dots, A[N]$.

The next Q lines of input will contain an operation, either in the form $1\ l\ r\ x$ for an operation of type 1 or $2\ l\ r$ for an operation of type 2.

Constraints

For all subtasks:

$$0 \leq A[i] \leq 10^5 \text{ for all valid } i.$$

$$1 \leq l \leq r \leq N$$

$$1 \leq x \leq 10^5$$

Subtask 1 [15%]

$$M = 2$$

$$1 \leq N, Q \leq 1\,000$$

Subtask 2 [15%]

$$M = 2$$

$$1 \leq N, Q \leq 100\,000$$

Subtask 3 [15%]

$$M = 3$$

$$1 \leq N, Q \leq 100\,000$$

Subtask 4 [15%]

$$M = 5$$

$$1 \leq N, Q \leq 100\,000$$

Subtask 5 [40%]

$$M = 10\,007$$

$$1 \leq N, Q \leq 200\,000$$

Output Specification

For each operation of type 2, output the answer on a new line.

Sample Input

```
2 5 3
1 2 3 4 5
2 1 4
1 2 5 7
2 1 5
```

Sample Output

```
0
1
```

Explanation

For the first operation, $1^2 + 2^2 + 3^2 + 4^2 = 30$, and $30 \equiv 0 \pmod{2}$.

For the second operation, the array A is now $\{1, 9, 10, 11, 12\}$.

For the third operation, $1^2 + 9^2 + 10^2 + 11^2 + 12^2 = 447$ and $447 \equiv 1 \pmod{2}$.