

# TLE '17 Contest 7 P3 - Countless Calculator Computations

**Time limit:** 1.0s    **Memory limit:** 256M  
Java: 2.0s  
Python: 3.0s

Leon likes to play with calculators whenever he gets bored in class. Such fascinating devices! One day, an intriguing problem occurred to him:

*If the equation  $X^{X^{X^{\dots}}} = Z$  contains  $Y$  number of  $X$ 's, then given the values of  $Y$  and  $Z$ , what is the approximate value of  $X$ ?*

Leon is tasked with  $Q$  queries regarding this problem. Apparently, he didn't perform these *countless calculator computations* well enough.

Can you help him?



*Leon is using a very powerful calculator.*

## Input Specification

The first line contains integer  $Q$  ( $1 \leq Q \leq 20\,000$ ), the number of queries.

The following  $Q$  lines each contain two space-separated integers  $Y_i$  ( $2 \leq Y_i \leq 100$ ) and  $Z_i$  ( $1 \leq Z_i \leq 2^{31} - 1$ ).

## Output Specification

For each query, output the approximate value of  $X_i$  on its own line, accurate within an absolute error of  $10^{-5}$ .

For 50% of the points,  $X_i$  may be accurate within an absolute error of  $10^{-1}$ .

## Sample Input

```
3
100 2
50 14
3 16
```

## Sample Output

```
1.414213562
1.4484039
1.99999
```